

# IOWA STATE UNIVERSITY

## Digital Repository

---

Integrated Crop Management News

Agriculture and Natural Resources

---

6-4-2013

## Sudden Death Syndrome and Heavy Spring Rains: Another Bad Year?

Daren S. Mueller

*Iowa State University*, [dsmuelle@iastate.edu](mailto:dsmuelle@iastate.edu)

Leonor F. S. Leandro

*Iowa State University*, [lleandro@iastate.edu](mailto:lleandro@iastate.edu)

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

---

### Recommended Citation

Mueller, Daren S. and Leandro, Leonor F. S., "Sudden Death Syndrome and Heavy Spring Rains: Another Bad Year?" (2013).  
*Integrated Crop Management News*. Paper 98.  
<http://lib.dr.iastate.edu/cropnews/98>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

[ICM Home](#)[ISU Extension Calendar](#)[Publications](#)[Extension News](#)[County Offices](#)[Contact Us](#)

search

Subscribe to Crop  
News


## Archives


[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)


## ISU Crop Resources


[Extension Field  
Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator  
Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology  
Extension](#)[Ag and Biosystems  
Engineering Extension](#)[Agribusiness  
Education Program](#)[Iowa Grain Quality  
Initiative](#)[College of Agriculture  
and Life Sciences](#)[ISU Extension](#)


# Integrated Crop Management NEWS

 PRINT STORY

 EMAIL STORY

 ADD TO DELICIOUS

 ATOM FEED

 FOLLOW ON TWITTER

## Sudden Death Syndrome and Heavy Spring Rains: Another Bad Year?

By Daren Mueller and Leonor Leandro, Department of Plant Pathology and Microbiology

Soybean sudden death syndrome (SDS) is one of the most damaging diseases of soybeans in Iowa and much of the Midwest. There are two phases of this disease – a root rot phase and a foliar symptom phase. SDS will be most problematic when weather conditions are conducive for disease development during both phases. The early cool, wet weather we have seen so far in 2013 helps increase the root rot phase of the disease. This can lead to development of severe SDS later in the growing season, as was seen in 2010 in Iowa.

In a recently published journal article, several plant pathologists at Iowa State University looked at rainfall, soil moisture and soil temperature in years with SDS (e.g., 2010) vs. years with little SDS (e.g., 2011). In this study, rainfall in April and May was similar in “SDS years” to “non-SDS years.” However, rainfall in June and July differed between disease years and non-disease years (Table 1). This highlights the importance of rainfall a bit later in the season to trigger the second phase of the disease. Also, soil temperature was less correlated to SDS severity compared to rainfall. The entire article is available in the [Plant Health Progress Journal on the Plant Management Network](#).

**Table 1. Average total precipitation in four years with high SDS prevalence (1993, 1998, 2008, and 2010) and five years with low SDS prevalence (2001, 2004, 2005, 2007, and 2011). Values are means of two locations: Ames (central Iowa) and Mount Pleasant (southeastern Iowa).**

|                         | Total Precipitation (cm) |      |      |      |        |
|-------------------------|--------------------------|------|------|------|--------|
| Year                    | April                    | May  | June | July | August |
| Mean SDS year           | 10.4                     | 15.1 | 26.3 | 19.5 | 14.7   |
| Mean non-SDS year       | 8.9                      | 15.4 | 11.5 | 8.9  | 12.2   |
| 30-yr Mean <sup>x</sup> | 9.0                      | 12.2 | 12.4 | 12.2 | 11.2   |

<sup>x</sup> Mean for the period of 1981 to 2010 at Ames and Mount Pleasant

One bit of good news for the 2013 season is that severe SDS usually is associated with early planting of soybeans. As many farmers are experiencing, the wet spring weather has delayed planting throughout the state. Fields with delayed planting should have less SDS develop in them. However, we have found research plots planted as late as June 15 still get SDS, so you are not completely out of the woods. SDS severity in these late

planted fields is very low, and much lower than fields planted in May.

*Daren Mueller is an assistant professor in the Department of Plant Pathology and Microbiology. He can be reached at 515-460-8000 or e-mail [dsmuelle@iastate.edu](mailto:dsmuelle@iastate.edu). Leonor Leandro is an Associate professor in the Department of Plant Pathology and Microbiology.*

---

This article was published originally on 6/4/2013. The information contained within the article may or may not be up to date depending on when you are accessing the information.

---

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.

---

Copyright ©2014 [Iowa State University Extension](#) | [Iowa State University](#)  
[Contact us](#) | [For Staff](#) | [Nondiscrimination and Information Disclosures](#) | [CMS Admin](#)  
Last Updated 6/4/2013